

3. *C. Nathorstii* Schuster. — Foliola lineari-lanceolata, sensim acuminata, basi contracta, 23–27 cm longa, 9–12 mm lata, pagina inferiore nitescens, subtus pallidiora et opaca, nervo utrinque prominente. Microsporophylla cuneata, 3–4 cm longa, 15–22 mm lata, area sterilis deltoidea, subito in acumen crassiusculum uncinatum eleganter arcuatim recurvatum, 1 cm longum attenuata. Macrosporophylla 15–30 cm longa, ovula 3–6 utrinque gerentia, lamina terminalis sterilis maximam latitudinem paululum supra basin attingens, basi paulum contracta, subtrigono-lanceolata vel subrhomboideo-lanceolata, elongata, 4–6,5 cm longa, 18–25 mm lata, in acumen longum excurrente, margine usque ad basin numerosis dentibus brevibus cristato-serrata, ovula margini macrosporophylli insidentia.

Ceylon: Thwaites 1866 n. 3689 in Herb. Barbey-Boissier. — Kultiviert Hort. Bogor. n. 38 (♂), n. 7 (folia). — Einh. Name: Madu. — Fig. 10 E.

Nota. Dubium mihi non est, quin hybrida ex *C. Rumphii* × *C. circinnalis* orta sit.