

Encephalartos imbricans (Zamiaceae): A New Species from Uganda

Author(s): P. Vorster and P. Heibloem

Source: Novon, Vol. 5, No. 4 (Winter, 1995), pp. 388-394

Published by: Missouri Botanical Garden Press Stable URL: http://www.jstor.org/stable/3391971

Accessed: 01/09/2008 09:56

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at http://www.jstor.org/action/showPublisher?publisherCode=mobot.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.

Encephalartos imbricans (Zamiaceae): A New Species from Uganda

P. Vorster and P. Heibloem

Department of Botany, University of Stellenbosch, Private Bag X1, 7602 Matieland, South Africa. Current address of P. Heibloem: P.O. Box 700, Nambour, Queensland 4560, Australia

ABSTRACT. Encephalartos imbricans, a new species from Uganda, is described, illustrated, and circumscribed in terms of other tropical African arborescent species. It is a large arborescent plant, with leaflets mostly strongly incubously overlapping. The strobili are green and glabrous; the megastrobilus is ovoid, shortly pedunculate, with the surfaces of the bullae wrinkled; the microstrobili emerge in succession, being narrowly ovoid, long-pedunculate, with the surfaces of the bullae smooth.

Encephalartos is widespread in the eastern parts of Africa, from the Eastern Cape Province of South Africa to southern Sudan, extending westward through equatorial Africa to Ghana and Nigeria (Goode, 1989: 244; Fig. 1). Nowhere is it common; it occurs mostly as relic populations, of which a considerable percentage must be considered to be in danger of extinction. The description of E. imbricans brings the number of described and recognized taxa to 57.

Encephalartos imbricans Vorster, sp. nov. TYPE: Uganda. Near northern shore of Lake Victoria, Oct. 1994, *P. Heibloem sub P. Vorster 3034* (holotype, K). Figures 2–5.

Planta magna arborescens. Foliola plerumque valde incuba, minime leviter falcata, dentibus in marginibus ambobus ad spationibus 25–50 mm et dentibus 3–4 confertis concentratis ad basin marginis superioris. Strobili virides, glabri, bullarum porcis mediolateralibus integris; megastrobili ovoidei, breve pedunculati, bullarum faciebus radialiter rugosis; microstrobili in successione emergentes, anguste ovoidei, longe pedunculati, bullarum faciebus laevigatis.

Large arborescent, unbranched plant; suckering profusely from base. Stem up to 6 m long, prostrate after 1.5–2.0 m from base, 400–650 mm diam., covered with old leaf bases. Fronds numerous, spreading, straight, emerging fresh medium green and maturing to dark glossy green, about 2(–3.5) m long, leaflets gradually reduced to prickles toward proximal part of petiole to leave no prickle-free petiole; median leaflets orientated

at angle of about 45° with rachis toward apex of frond, opposing leaflets are in the same plane as seen in cross section of leaf, very narrowly ovate and very slightly falcate, with teeth 25-50 mm apart on both margins and often 3-4 closely spaced teeth near base of upper margin, mostly strongly incubously overlapping, 200-250 mm long and up to 22 mm wide. Megastrobili 2-3, ovoid with apical sporophylls sterile, shortly pedunculate with exposed peduncle 30-40 mm long, glaucous green, glabrous, about 300 mm long and 180 mm diam.; bullae of sporophylls moderately projecting, terminal facet centrally situated and about 1/3 the horizontal diameter of bulla, median facet differentiated in most sporophylls, ridges separating facets clearly differentiated and simple, terminal facet smooth but other facets radially rugose. Microstrobili 5-8, emerging and maturing in succession, each carried on a stout peduncle up to 300 mm long; strobili glaucous green, glabrous, about 300 mm long and 90 mm diam. at maturity; bullae of sporophylls moderately projected, slightly drooping, only terminal facet differentiated and about 1/3 the horizontal diameter of bulla, smooth. Seed covered with orange (not red) sarcotesta, about 45-50 mm long and 25-30 mm wide; with sarcotesta removed about 40 mm long and 18-24 mm wide.

Variation. A concentration of 3 to 4 closely spaced teeth near the base of the upper margin is present in most (Fig. 2B), but not all (Fig. 3), leaflets. Likewise, the majority of plants have strongly overlapping leaflets; but through a quirk of fate the frond collected for the type specimen has well-spaced leaflets—it probably developed in a situation where it was shaded by other plants and thus became etiolated.

Phenology. During October the plants had fresh male and female cones, and seeds from the previous season were scattered about. A mature megastrobilus collected at this time disintegrated in mid-November, suggesting that the megastrobili take slightly more than a year to mature.

Geographical distribution and habitat. Known




Figure 1. Encephalartos imbricans Vorster. Known geographical distribution in relation to distributions of other tropical African arborescent species. Based on Stevenson et al. (1990b) and Goode (1989). Scale bar = 1000 km.

only from two populations on the northern shore of Lake Victoria, where it occurs on two adjacent low hills in otherwise fairly flat country (Fig. 1). It grows on bare granite outcrops, wedged in crevices from which the plants cannot escape, in minimal soil, and under intense insolation. These outcrops (Fig. 6) have minimal vegetation, yet the charred trunks of *Encephalartos imbricans* testify that the plants are subject to periodic burning, and suitable niches for seedling establishment must be very scarce.

Conservation status. The population visited consisted of about 300 individuals. It appeared to be vigorous and was coning, yet no young plants or

evidence of seedling regeneration were seen. It seems likely that the number of plants is in equilibrium with the limited extent of the specialized habitat. Seed dissected contained no embryos, and no insects that could be suspected of being pollen vectors were found in either male or female strobili. We therefore conclude that these populations are vulnerable to overcollecting, and, accordingly, the precise locality is not made public.

Affinities. Within Encephalartos, interspecific relationships are still poorly understood. For many years species were separated solely on vegetative characteristics (Vorster, 1993). There does indeed seem to be a considerable amount of convergence

390 Novon




Figure 2. Encephalartos imbricans Vorster. —A. Proximal part of petiole. —B. Median leaflet. —C. Megasporophyll in ventral view. —D. Seed with sarcotesta. —E. Seed with sarcotesta removed. —F. Microsporophyll in ventral view. Scale bar = 50 mm. (Heibloem sub Vorster 3034.)

Figure 3. Encephalartos imbricans Vorster. Part of median zone of frond, showing incubously overlapping leaflets. Scale bar = 100 mm. (Plant in habitat.)

in cone characteristics, yet these are often useful to circumscribe species, if not always reflecting phylogenetic relationships. It would appear that recurring vegetative characteristics may well be cryptic signposts of evolutionary kinship. The following is therefore intended to identify *E. imbricans*, but is certainly not intended to reflect phylogenetic relationships.

Encephalartos imbricans resembles E. bubalinus Melville (Melville, 1957) in its green, ovoid, shortly pedunculate megastrobili (Fig. 4E), but differs by the somewhat wrinkled instead of smooth facets of the megastrobilate bullae (Fig. 4F), and by the long-peduncled instead of almost sessile microstrobili. In both species the leaflets are overlapping, but in E. imbricans the leaflets taper gradually to their apices (Fig. 2B), whereas in E. bubalinus they are uncinate.

Encephalartos imbricans resembles an undescribed species from western Uganda in its green, ovoid megastrobili, but is readily distinguished by the somewhat wrinkled instead of smooth facets of the megastrobilate bullae, and the mostly strongly overlapping (Fig. 3) instead of well-spaced leaflets. The concentration of 3—4 teeth near the base of the upper leaflet margin was not seen in the western Ugandan species.

Encephalartos imbricans resembles E. laurentianus De Wildeman (De Wildeman, 1903a, b, c, 1907; Bois, 1907; Gentil, 1904a, b; Lebrun, 1930; Robyns, 1948) by its shortly pedunculate, ovoid megastrobili, of which the facets of the bullae are somewhat wrinkled, as well as by the large number of long-pedunculate microstrobili; it is distinguished by the absence of a characteristic thin indumentum of reddish brown trichomes, and green rather than greenish yellow pigmentation of the strobili. Vegetatively, E. laurentianus is a very different species, with much larger (up to 500 mm long and 50 mm wide) and softer-textured leaflets.

The new species resembles Encephalartos septentrionalis Schweinfurth (Goode, 1989: 236) by its pedunculate, ovoid megastrobili, and long-pedunculate microstrobili, but differs in the very short instead of very long (up to 300 mm) peduncled megastrobili (Goode, 1989: 239) and the glaucous green instead of greenish yellow pigmentation of the strobili. Both species tend to have a concentration of up to 4 teeth near the base of the upper margin of the leaflet, but in E. septentrionalis the leaflets are not overlapping but well spaced. Despite being an old species, E. septentrionalis is not well known, and in view of its unusually wide geographical distribution (according to the map in Stevenson et al., 1990b: 157) more than one species may be involved. In this respect it is worth noting that the very long megastrobilate peduncle depicted by Goode (1989: 239) is not mentioned elsewhere in the literature.

Encephalartos imbricans resembles E. sclavoi De Luca, Stevenson & Moretti (Stevenson et al., 1990a, b) by the ovoid, shortly pedunculate megastrobili as well as by the overlapping leaflets; it differs in that both mega- and microstrobili are green with centrally placed terminal facets on rugose megastrobilate bullae instead of orange-yellow with terminal facets of the smooth megastrobilate bullae situated toward the proximal margins of the bullae, the long (up to 300 mm) instead of very short-peduncled (20–40 mm) microstrobili, and the narrower (up to 22 mm vs. up to 40 mm) leaflets. Encephalartos sclavoi lacks the concentration of 3 to 4 teeth near the base of the upper leaflet margin.

Both Encephalartos imbricans and E. turneri Lavranos & Goode (Lavranos & Goode, 1985) have imbricate leaflets, but in E. turneri they are succubous instead of incubous (Fig. 3), they are wider

392 Novon




Figure 4. Encephalartos imbricans Vorster. —A. Microstrobilus in immature stage. —B. Portion of immature microstrobilus, showing centrally situated terminal facet. —C. Microstrobilus at anthesis. —D. Portion of microstrobilus at anthesis, showing drooping terminal facet. —E. Megastrobilus. —F. Portion of megastrobilus showing rugose facets of bullae. Scale bars (at the right in each photograph): A, C, E = 100 mm; B, D, F = 10 mm. (Heibloem sub Vorster 3034.)



Figure 5. Encephalartos imbricans Vorster. Plant in habitat, showing habit. Scale bar = 1 m.

(up to 30 mm vs. up to 22 mm), and characteristically the apices are uncinate instead of tapering to a fine point. In *E. turneri* the megastrobilus is narrowly ovoid instead of ovoid, completely sessile instead of shortly peduncled, and the bullae project more. It bears fewer microstrobili (up to 3 vs. 5–8) on shorter peduncles (up to 120 vs. to 300 mm). In both sexes the cones are not green but yellowish with a conspicuous pink bloom (Lavranos & Goode, 1985).

Encephalartos imbricans resembles E. ituriensis Bamps & Lisowski (Bamps & Lisowski, 1990) by its long-peduncled microstrobili. However, in E. ituriensis the megastrobilus is shortly cylindrical and the strobili of both sexes are yellow instead of green (Goode, 1989). Moreover, the type collection of E. ituriensis (Lisowski 41057, BR) leaves the impression of relatively soft textured and flaccid leaflets as in E. laurentianus, in marked contrast to the hard and rigid leaflets of E. imbricans.

Other east African arborescent species with which we compared *Encephalartos imbricans* are *E. hildebrandtii* A. Braun & Bouché (Prain, 1915;

Stapf, 1914), E. kisambo Faden & Beentje (Faden & Beentje, 1989; Moretti et al., 1989), and E. tegulaneus Melville (Melville, 1957). From all these it differs by its glaucous green instead of yellow cones, and the ovoid instead of cylindrical megastrobili. Yet in various respects it resembles these species, which probably indicates a near relationship. In all four species the microstrobili are borne on very long peduncles, which can be up to 300 mm long. In both E. imbricans and E. hildebrandtii the microstrobili emerge and mature in succession rather than simultaneously. This trait is shared by the undescribed species from western Uganda, by E. gratus Prain (Prain, 1916) from Malawi, and E. ferox G. Bertoloni (pers. obs.) from southern Mozambique and South Africa, and is suspected to be present in at least some other tropical African species including E. septentrionalis and E. laurentianus. In the South African species, in the E. manikensis (Gilliland) Gilliland complex, the E. poggei Ascherson & Graebner group, and in E. barteri Carruthers, the cones emerge simultaneously (teste Goode, 1989; pers. obs.). The conspicuous grouping of 3 to 4 teeth near the base of the upper leaflet margin is shared by E. hildebrandtii (some specimens, e.g., Faden, Faden & Rathbun 72 136, K), E. kisambo, E. tegulaneus, E. bubalinus (where it is poorly expressed), the Malawian E. gratus, and the South African E. woodii Sander, and it is tempting to read phylogenetic significance in it.

Melville (1957: 248) briefly described a variety dentatus of E. hildebrandtii, based on cultivated specimens. Heenan (1977: 281–282) reported that he had rediscovered it near Jinja in Uganda. This is not the same as E. imbricans, because E. hildebrandtii var. dentatus is distinguished by divaricate-dentate medio-lateral ridges on both the micro- and megasporophyllate bullae.

Acknowledgments. The research of which this paper is a result, was financed by the University of Stellenbosch and the South African Foundation for Research Development. We thank the Keeper of Botany at K for granting to P. V. the opportunity to examine material of Encephalartos in K, and for providing us with photocopies of rare literature. For assistance with the latter we also wish to record our appreciation to the South African liaison officer at Kew, Miss M. Koekemoer. P. H. wishes to express his gratitude to Paul C. Mackenzie of Nairobi, Kenya, for logistic and other support during the course of his fieldwork.

394 Novon



Figure 6. Encephalartos imbricans Vorster. Plants in typical habitat, wedged in crevices in granite outcrop. Scale bar = 1 m.

Literature Cited

Bamps, P. & S. Lisowski. 1990. A new species of Encephalartos [E. ituriensis] (Zamiaceae) from northeastern Zaire. Mem. New York Bot. Gard. 57: 152–155.

Bois, D. 1907. Les Encephalartos lemarinelianus et laurentianus. Rev. Hort. 79: 177–179.

Faden, R. B. & H. J. Beentje. 1989. Encephalartos kisambo, a new cycad from Kenya, with a note on E. tegulaneus. Utafiti 2: 7-10.

Gentil, L. 1904a. Encephalartos from the Congo. Gard. Chron., June 1904: 370–371.

——. 1904b. Les cycadées au Congo. Rev. Hort. Belge étrangère 30: 7–9.

Goode, D. 1989. Cycads of Africa. Struik Winchester, Cape Town.

Heenan, D. 1977. Some observations on the cycads of central Africa. J. Linn. Soc., Bot. 74: 277-288.

Lavranos, J. & D. Goode. 1985. Encephalartos turneri (Cycadaceae), a new species from Mozambique. Garcia de Orta 7: 11-14.

Lebrun, J. 1930. Note sur l'Encephalartos laurentianus De Wild., de la vallée du Kwango. Rev. Zool. Bot. Africaines 19: 384–390.

Melville, R. 1957. *Encephalartos* in central Africa. Kew Bull. 2: 237–257.

Moretti, A., P. de Luca, J. P. Sclavo & D. W. Stevenson. 1989. *Encephalartos voiensis* (Zamiaceae), a new east central African species in the *E. hildebrandtii* complex. Ann. Missouri Bot. Gard. 76: 934–938.

Prain, D. 1915. Encephalartos hildebrandtii. Bot. Mag. 141: tt. 8592 & 8593.

1916. Diagnoses Africanae LXVII. Kew Bull. 7: 180-182.

Robyns, W. 1948. Cycadaceae. Pp. 1–4 in Flore du Congo belge et du Ruanda-Urundi, Spermatophytes, vol. 1. I.N.É.A.C., Bruxelles.

Stapf, O. 1914. Encephalartos hildebrandtii. Kew Bull. 10: 386–392.

Stevenson, D. W., A. Moretti & P. de Luca. 1990a. Encephalartos sclavoi De Luca, D. Stevenson & Moretti (Zamiaceae), a new species from Tanzania. Delpinoa, n.s. 29–30: 3–5.

——, —— & ——. 1990b. A new species of Encephalartos [E. sclavoi] (Zamiaceae) from Tanzania. Mem. New York Bot. Gard. 57: 156–161.

Vorster, P. 1993. Taxonomy of Encephalartos (Zamiaceae): Taxonomically useful external characteristics. Pp. 294–299 in D. W. Stevenson & K. J. Norstog (editors), Proceedings of CYCAD 90, the Second International Conference on Cycad Biology, Palm & Cycad Societies of Australia, Milton, Queensland.

De Wildeman, É. 1903a. Études de systématique et de géographie botaniques sur la Flore du Bas- et du Moyen-Congo. Ann. Mus. Congo, Sér. 5, Bot. 1: 9–10, tt. 23–25.

1903b. Icones Selectae Horti Thenensis 4: 183.
1903c. Notices sur des plantes utiles ou intéressantes de la Flore du Congo [1]: 386-394.

Vanbuggenhout, Bruxelles.