

A SYNOPSIS
OF
THE QUEENSLAND FLORA;

CONTAINING BOTH THE
PHÆNOGAMOUS AND CRYPTOGAMOUS PLANTS.

BY
FREDK. MANSON BAILEY, F.L.S.,
ETC., ETC.,
COLONIAL BOTANIST,
Author of "Queensland Ferns," "The Fern World of Australia," &c., &c.

BRISBANE:
JAMES C. BEAL, GOVERNMENT PRINTER, WILLIAM STREET,
—
1883.

M. Douglasii, *W. Hill, inedit.* (After the Hon. John Douglas.) Stem short, rarely over 3 feet high above the surface, but, like many others of this genus, often reaching some distance beneath; diameter about 1 foot. Leaves somewhat channelled, 2 to 4 feet long, petioles about 1 foot; leaflets numerous, broad-linear, gradually narrowed to the pointed apex, about $\frac{1}{2}$ -inch broad at the widest part, very few of the leaflets abbreviated at the base of the leaf. Male cone about 7 inches long, diameter at widest part about 3 inches; stalk 1 to $1\frac{1}{2}$ feet long, glabrous; scales glabrous, the apex triangular, not spinulous. Female cones about 1 foot long and 5 inches broad, the apex of scales triangular, flat, spineless, $1\frac{1}{2}$ to 2 inches broad. Nuts orange, over 1 inch long. Native name, "Coobine." (*Fraser's Island.*)