

## Notes on the Genus *Macrozamia*.

By CHARLES MOORE, F.L.S., Vice-President of the Royal Society of N.S.W., Director of the Botanic Gardens, Sydney.

[Read before the Royal Society of N.S.W., 5 September, 1884.]

120

### NOTES ON THE GENUS *MACROZAMIA*.

6. *Macrozamia Fawcetti*, C. Moore.—Trunk and base of the petioles covered with a dense tomentum. Leaves varying from 2 to 4 feet in length, of a dark glossy green colour above, paler beneath. Glabrous in an adult state, hairy when young. Pinnae about three-quarters of an inch in breadth and from 6 to 7 inches in length, semi-falcate, broader upwards, and terminating with an abrupt falcate point, which is slightly but sharply toothed. Lower part of the rachis flat on the upper surface, rather keel-shaped below, gradually rounding in form

### NOTES ON THE GENUS *MACROZAMIA*.

121

upwards till it becomes almost terete towards the apex. Male cones on rather long peduncles, woolly at the base, above 8 inches long and 2½ inches broad; scales flat, terminating abruptly with sharp points. Fruiting cones not seen.

*Habitat*.—On high ground on the upper part of the Richmond; discovered by C. Fawcett, Esq., P.M. Very little is known of the habit of this species. Only freshly-gathered leaves and old male cones have as yet been seen, but this proves that it is very different from any species hitherto described.