

1

АСТА
HORTI PETROPOLITANI.

TOMUS IV.
Fasciculus II.

ТРУДЫ
ИМПЕРАТОРСКАГО
С.-ПЕТЕРБУРГСКАГО БОТАНИЧЕСКАГО САДА.

ТОМЪ IV.
ВЫПУСКЪ II.



longa, 25—35 nervia, usque 8 c. m. longa et $1\frac{1}{2}$ — $2\frac{1}{2}$ c.
lata.

Habitat in regno mexicano.

9. *Zamia manicata* Linden.

Linden in horto Lindeano.

Truncus cylindricus. Petioli sparsim aculeati. Folia pauci-plurijugopinnata. Foliola elliptica, acuminata, undulata, apicem versus obsolete crenulato-denticulata, basin versus in petiolum annulo prominente manicatum attenuata.

10. *Zamia boliviana* D. C.

D. C. prodr. XVI. II. 540. — *Ceratozamia boliviana* Brongn. in Ann. sc. nat. ser. III. tom. V. p. 9. — Z. Brongniarti Wedd. in Ann. sc. nat. ser. III. tom. XIII. pag. 249 tab. 4. — Eichl. in Mart. fl. bras. tom. IV. pag. 413 tab. 108.

Truncus hypogaeus, cylindraceus. Perulae e basi latiore lanceolatae, sensim attenuato-acutae. Folia 35—60 c. m. longa. Petiolus inermis, 15—35 c. m. longus, teres, juxta axillam tomentosus. Rhachis obtuse triquetra, supra leviter carinata, circiter petioli longitudine. Foliola 10—18 juga, anguste lineari-lanceolata, subfalcata, circiter 12 nervia, sensim in apicem acutum attenuata, integerrima v. apice utrinque obsolete 3—5 denticulata, 15—25 c. m. longa, circiter 8 m. m. lata. Strobili pedunculati, cylindracei; masculi 7—9 c. m. longi; foeminei quam masculi subduplo longiores et quadruplo crassiores, ferrugineo-virescentes, apice acuminati.

Habitat in Bolivia inque Brasilia.

Vix ac ne vix quidem a sequenti specie differre videtur.

11. *Zamia media* Jacq. (v. v.)

Jacq. hort. Schönbr. tab. 397 et 398. — Miq. monogr. pag. 68. — D. C. prodr. XVI. II. 542. — Z. integrifolia Rich. Conif. et Cyc. pag. 191 tab. 27 et 28.